

On amorphic C -algebras

I.N. Ponomarenko¹

Petersburg Department of V.A. Steklov
Institute of Mathematics
Fontanka 27, St. Petersburg 191023, Russia
inp@pdmi.ras.ru
<http://www.pdmi.ras.ru/~inp>

A. Rahnamai Barghi

Institute for Advanced Studies in Basic Sciences
Gava Zang, P.O.Box: 45195-1159 Zanjan, Iran

email: rahnama@iasbs.ac.ir

¹Partially supported by RFFI, grants 03-01-000349, 05-01-00-899, NSH-2251.2003.1.
The author thanks the Institute for Advanced Studies in Basic Sciences (IASBS) for its hospitality during the time that this paper was written

running title: On amorphic C -algebras

SEND PROOFS TO:

A. Rahnamai Barghi

Institute for Advanced Studies in Basic Sciences

Gava Zang, P.O.Box: 45195-1159 Zanzan, Iran

email: rahnama@iasbs.ac.ir

ABSTRACT

An amorphic association scheme has the property that any of its fusion is also an association scheme. In this paper we generalize the property to be amorphic to an arbitrary C -algebra and prove that any amorphic C -algebra is determined up to isomorphism by the multiset of its degrees and an additional integer equal ± 1 . Moreover, we show that any amorphic C -algebra with rational structure constants is the fusion of an amorphic homogeneous C -algebra. Finally, as a special case of our results we obtain the well-known Ivanov's characterization of intersection numbers of amorphic association schemes.

AMS Subject Classification 2000: 20C05 05E30

Key words and phrases: C-algebra, association scheme, table algebra

1 Introduction

One of the important problems in association schemes theory consists of the constructive enumeration of all the subschemes of a given scheme. Such a problem becomes intractable when the number of subschemes is large. In the limit case this number coincides with the number of all partitions of the set of classes of the scheme in question. In order to describe this situation, the notion of amorphic cellular ring (the adjacency ring of an association scheme) was introduced in [10]. In paper [12] that is a continuation of [10], a complete characterization of the intersection numbers array of an arbitrary amorphic cellular ring is given. Since then a number of examples of amorphic association schemes were found (see [6, 13]). However, the problem of characterization of amorphic association schemes is far from being solved.

The notion of C -algebra goes back to the old paper of Kawada (1942) in which a finite dimensional analog of the Tannaka duality was introduced (see Section 2 in the below for a background of C -algebras). The two basic examples of C -algebras (which are also table algebras in sense of [1]) are the algebra of conjugacy classes of a finite group and the character algebra of a finite group. Besides, in [4] it was showed that the adjacency algebra of a commutative association scheme is a C -algebra. A special case of non-commutative C -algebras (including all the above examples and also the

adjacency algebra of an arbitrary association scheme) was studied in [2].

The main purpose of the paper is to generalize the concept of amorphic association scheme for C -algebras and study amorphic C -algebras in details. According to our definition (see Definition 3.1), the adjacency algebra of an amorphic association scheme is an amorphic C -algebra, but not all of the latter arise in this way. For instance, affine C -algebras corresponding to affine planes of order q (see Section 2) come from affine association schemes and so are amorphic due to [10]. On the other hand, there are some generalized affine C -algebras (see Subsection 3.3) which do not come from association schemes.

After studying some elementary properties of amorphic C -algebras, we prove that each such algebra is determined up to isomorphism by the multiset of its degrees and an additional integer equal ± 1 (Theorem 3.3). As a special case of this result we obtain the well-known Ivanov's characterization of intersection numbers of amorphic association schemes [12]. In Section 4, the class of homogeneous amorphic C -algebras is studied (this class contains generalized affine C -algebras mentioned in the above). We show that any nontrivial amorphic C -algebra with integer structure constants is a fusion of a generalized affine algebra (Corollary 4.2). Furthermore, we give a necessary and sufficient condition for a C -algebra to be homogeneous and amorphic (Theorem 4.3). Finally it is proved that the dual of a homogeneous amorphic table algebra is a homogeneous amorphic table algebra too (Theorem 4.4).

Concerning association schemes and permutation groups we refer the reader to [4] and [8] respectively.

2 C -algebras

In the definition below we follow the paper [9] in which arbitrary non-commutative C -algebras were studied from the duality theory point of view.

Let A be an associative algebra over $\mathbb{C}$ with the unit e , semilinear involution anti-automorphism $*$ and a linear basis $\mathcal{R}$. The structure constants of this algebra with respect to the basis $\mathcal{R}$ are defined by

$$r \cdot s = \sum_{t \in \mathcal{R}} c_{r,s}^t t, \quad r, s \in \mathcal{R}. \quad (1)$$

Definition 2.1 The pair $\mathcal{A} = (A, \mathcal{R})$ is called a *C-algebra*, if the following conditions are satisfied:

- (C1) $e \in \mathcal{R}$ and $r^* \in \mathcal{R}$ for all $r \in \mathcal{R}$,
- (C2) $c_{r,s}^t \in \mathbb{R}$ for all $r, s, t \in \mathcal{R}$,
- (C3) for all $r, s \in \mathcal{R}$ the equality $c_{r,s}^e = d_r \delta_{r^*,s}$ holds where $d_r > 0$ and δ is the Kronecker symbol,
- (C4) a linear functional on A , defined by the mapping $r \mapsto d_r$, $r \in \mathcal{R}$, is a one-dimensional $*$ -representation of the algebra A .

The *C-algebra* $\mathcal{A}$ is called *symmetric* (resp. *commutative*) if the involution $*$ is trivial (resp. the algebra $\mathcal{A}$ is commutative). Obviously, any symmetric *C-algebra* is commutative.

Two *C-algebras* $\mathcal{A} = (A, \mathcal{R})$ and $\mathcal{A}' = (A', \mathcal{R}')$ are called *isomorphic*, if there exists an $*$ -algebra isomorphism $\varphi : A \rightarrow A'$ such that $\varphi(\mathcal{R}) = \mathcal{R}'$; such an isomorphism is called the *isomorphism* from $\mathcal{A}$ to $\mathcal{A}'$. For $\mathcal{A} = \mathcal{A}'$ the set of all of them form a group $\text{Aut}(\mathcal{A})$ called the *automorphism group* of the *C-algebra* $\mathcal{A}$. From the definition it follows that this group acts faithfully on the set $\mathcal{R}$ and hence it can be naturally identified with a subgroup of the group $\text{Sym}(\mathcal{R}^\#)$ where $\mathcal{R}^\# = \mathcal{R} \setminus \{e\}$.

Example: affine C-algebras (cfg. [10]). An *affine plane* is a triple consisting of a set P of points, a set L of lines and an incidence relation between the points and the lines such that

- (A1) any two different points are incident to a unique line,
- (A2) for any point p and for any line l not incident to p there exists a unique line incident to p and parallel to l ,
- (A3) there exist three different points not incident to the same line.

From now on we assume that the plane is finite, i.e. the set $P \cup L$ is finite. Then there exists an integer $q \geq 2$, called the *order* of the plane, such that $|P| = q^2$ and $|L| = q^2 + q$, and each line incident to exactly $q + 1$ points. Besides, there are exactly $q + 1$ classes $B_1, \dots, B_{q+1}$ of pairwise parallel (non-intersecting) lines, each of which is of cardinality q .

Let Mat_P be the full matrix algebra of degree $|P|$ over $\mathbb{C}$ whose rows and columns are indexed by the elements of P . Given $i = 1, \dots, q+1$ we define a $\{0,1\}$ -matrix r_i belongs to Mat_P by

$$(r_i)_{u,v} = \begin{cases} 1, & \text{if } u \neq v \text{ and } l(u,v) \in B_i, \\ 0, & \text{otherwise,} \end{cases}$$

where $l(u,v)$ is the line incident to both u and v . One can see that

$$r_i r_j = \begin{cases} (q-1)r_0 + (q-2)r_i, & \text{if } i = j, \\ \sum_{k \neq 0, i, j} r_k, & \text{if } i \neq j \end{cases} \quad (2)$$

where r_0 is the identity matrix of Mat_P . So the set $\mathcal{R} = \{r_i : i = 0, \dots, q+1\}$ is a linear base of the subalgebra A of the algebra Mat_P generated by $\mathcal{R}$. Since conditions (C1)–(C4) are trivially satisfied (with $*$ being the Hermitian conjugation in Mat_P), the pair $\mathcal{A} = (A, \mathcal{R})$ is a C -algebra. We call it the *affine C -algebra*. From (2), it follows that any affine C -algebra is determined up to isomorphism by the order of the corresponding affine plane (and does not depend on the choice of the plane). Moreover, it is easy to see that $\mathcal{A}$ is symmetric and $\text{Aut}(\mathcal{A}) \simeq \text{Sym}(q+1)$. ■

Let $\mathcal{A} = (A, \mathcal{R})$ be a C -algebra. Then the structure constants $c_{r,s}^t$ of the algebra A with respect to the base $\mathcal{R}$ satisfy some relations. For instance, from (C4) it follows that $d_e = 1$ and $d_r = d_{r^*}$ for all $r \in \mathcal{R}$. Besides, the associativity of A exactly means that given $r, s, t \in \mathcal{R}$ the following equalities hold:

$$\sum_{v \in \mathcal{R}} c_{r,s}^v c_{v,t}^u = \sum_{v \in \mathcal{R}} c_{r,v}^u c_{s,t}^v, \quad u \in \mathcal{R}. \quad (3)$$

In fact, the left-hand side and the right-hand side of the equalities (3) are the coefficients of u in the products $(rs)t$ and $r(st)$ respectively. Take $u = e$. Then from condition (C3) it follows that

$$d_t c_{r,s}^{t^*} = d_r c_{s,t}^{r^*} = d_s c_{t,r}^{s^*}, \quad r, s, t \in \mathcal{R}. \quad (4)$$

Finally, applying the linear functional from condition (C4) to all the products of two elements of $\mathcal{R}$, we get the equalities

$$d_r d_s = \sum_{t \in \mathcal{R}} d_t c_{r,s}^t, \quad r, s \in \mathcal{R}. \quad (5)$$

Below we denote by $D(\mathcal{A})$ the multiset $\{d_r : r \in \mathcal{R}\}$ and call the number $n(\mathcal{A}) = \sum_{r \in \mathcal{R}} d_r$ the *degree* of $\mathcal{A}$. In particular, if $\mathcal{A}$ is the affine C -algebra corresponding to an affine plane of order q , then from (2) it follows that $D(\mathcal{A}) = \{1, q-1, \dots, q-1\}$ ($q+1$ times) and $n(\mathcal{A}) = q^2$. This gives an example of a *homogeneous* C -algebra, i.e. one for which the number d_r does not depend on $r \in \mathcal{R}^\#$.

Let π be a partition of $\mathcal{R}$ such that $\{e\} \in \pi$. Then the pair $\mathcal{A}_\pi = (A_\pi, \mathcal{R}_\pi)$ where A_π is the linear subspace of A spanned by the set

$$\mathcal{R}_\pi = \left\{ \sum_{r \in R} r : R \in \pi \right\},$$

is called the π -*fusion* of $\mathcal{A}$. It is easy to see that $\mathcal{A}_\pi$ is a C -algebra iff A_π is an $*$ -subalgebra of A . For example, the latter is always true whenever $|\pi| \in \{2, |\mathcal{R}|\}$. In this paper we are interesting in the case when each fusion of $\mathcal{A}$ is a C -algebra.

3 A characterization of amorphic C -algebras

3.1 The following definition extends the notion of amorphic association scheme to arbitrary C -algebras.

Definition 3.1 *A C -algebra $\mathcal{A} = (A, \mathcal{R})$ is called amorphic if for each partition π of $\mathcal{R}$ the π -fusion $\mathcal{A}_\pi = (A_\pi, \mathcal{R}_\pi)$ of $\mathcal{A}$ is a C -algebra.*

From the definition it follows that the adjacency algebra of an amorphic association scheme is an amorphic C -algebra. Next, obviously, any C -algebra of dimension at most 3 is amorphic; we call them *trivial* amorphic C -algebras. Nontrivial examples of amorphic C -algebras can be obtained from [10, Theorem 3.3] which implies that any affine C -algebra is amorphic (see also paper [6] and the references in it). A full characterization of amorphic C -algebras isomorphic to the adjacency algebras of association schemes follows from [12] (a full characterization of amorphic cyclotomic association schemes over finite fields up to association scheme isomorphisms follows from [5]). However, there is a number of C -algebras which are not isomorphic to the adjacency algebra of an association scheme (e.g. the structure constants of the latter must be nonnegative integers) and our results show that such examples can be found in the class of amorphic C -algebras.

Let $\mathcal{A} = (A, \mathcal{R})$ be an amorphic C -algebra. Given $r, s \in \mathcal{R}^\#$ set $\pi = \pi_{r,s}$ to be the partition of $\mathcal{R}$ with classes $\{e\}$, $\{r\}$, $\{s\}$ and $\mathcal{R}^\# \setminus \{e, r, s\}$ (we don't assume that $r \neq s$). Since $\mathcal{A}_\pi = (A_\pi, \mathcal{R}_\pi)$ is a C -algebra with $r, s \in \mathcal{R}_\pi$, we see that

$$c_{r,s}^u = c_{r,s}^v, \quad u, v \in \mathcal{R}^\# \setminus \{r, s\}. \quad (6)$$

Moreover, for $r = s$ we also have $r^* \in \{r, t\}$ where t is the sum of all $u \in \mathcal{R}^\# \setminus \{r\}$. If $|\mathcal{R}| \geq 4$, this implies by condition (C1) that $r = r^*$. Thus any nontrivial amorphic C -algebra is symmetric and hence any amorphic C -algebra is commutative (c.f. [10, Theorem 2.3]).

Theorem 3.2 *Let $\mathcal{A} = (A, \mathcal{R})$ be a symmetric C -algebra. Then the following three statements are equivalent:*

- (1) $\mathcal{A}$ is amorphic,
- (2) given $r, s \in \mathcal{R}^\#$ the $\pi_{r,s}$ -fusion of $\mathcal{A}$ is a C -algebra,
- (3) given $r, s \in \mathcal{R}^\#$ the equalities (6) hold.

Proof. The implications (1) $\Rightarrow$ (2) and (2) $\Rightarrow$ (3) are obvious. To prove the implication (3) $\Rightarrow$ (1) let π be a partition of $\mathcal{R}$ with $\{e\} \in \pi$. Then the linear space A_π is closed with respect to $*$ because the C -algebra $\mathcal{A}$ is symmetric. Given $R, S \in \pi$ set $T = \mathcal{R} \setminus (R \cup S \cup \{e\})$. Then from (6) it follows that

$$\sum_{r \in R} \sum_{s \in S} c_{r,s}^u = \sum_{r \in R} \sum_{s \in S} c_{r,s}^v, \quad u, v \in T.$$

This implies that $\sum_{r \in R} r \sum_{s \in S} s \in A_\pi$ and so the linear space A_π is an $*$ -subalgebra of A . Thus the fusion $\mathcal{A}_\pi$ is a C -algebra and we are done. ■

3.2 The following theorem gives explicit formulas for the structure constants of a nontrivial amorphic C -algebra $(A, \mathcal{R})$. We observe that due to equalities (4) it suffices to find the structure constants of the form $c_{r,s}^t$ where $r, s, t \in \mathcal{R}^\#$ are such that $r \neq s \neq t \neq r$ or $r = s \neq t$ or $r = s = t$.

Theorem 3.3 *Let $\mathcal{A} = (A, \mathcal{R})$ be a nontrivial amorphic C -algebra. Then the structure constants of $\mathcal{A}$ are uniquely determined by the multiset $D(\mathcal{A})$ and by the number $\varepsilon \in \{-1, +1\}$. More exactly, given $r, s, t \in \mathcal{R}^\#$ we have*

$$c_{r,s}^t = \begin{cases} \frac{d_r d_s}{(\sqrt{n} + \varepsilon)^2}, & \text{if } r \neq s \neq t \neq r, \\ \frac{d_r}{2} \left(1 + \frac{d_r - d'_r}{(\sqrt{n} + \varepsilon)^2} \right), & \text{if } r = s \neq t, \\ -\frac{d'_r}{2} \left(1 + \frac{d_r - d'_r}{(\sqrt{n} + \varepsilon)^2} \right) + d_r - 1, & \text{if } r = s = t, \end{cases} \quad (7)$$

where $n = n(\mathcal{A})$ and $d'_r = n - 1 - d_r$.

Proof. Given two distinct elements $r, s \in \mathcal{R}^\#$ set $\pi = \pi_{r,s}$ and t to be the sum of all $u \in \mathcal{R}^\# \setminus \{r, s\}$. Then $\mathcal{A}_\pi = (A_\pi, \mathcal{R}_\pi)$ is an amorphic C -algebra of dimension 4 with $\mathcal{R}_\pi = \{e, r, s, t\}$. Since it is symmetric, formulas (5) and (6) imply that

$$\begin{cases} d_r^2 - d_r &= d_r c_{r,r}^r + d_s c_{r,r}^s + d_t c_{r,r}^t \\ d_r d_s &= d_r c_{r,s}^r + d_s c_{r,s}^s + d_t c_{r,s}^t \\ 0 &= d_t c_{r,r}^s - d_s c_{r,r}^t \end{cases} \quad (8)$$

where $d_t = n - 1 - d_r - d_s$. By (4) the number $d_w c_{u,v}^w$ does not depend on $u, v, w \in \mathcal{R}_\pi^\#$ such that $\{u, v, w\} = \{r, s, t\}$. Denote it by T and set $X_{uv} = d_v c_{u,u}^v$ where $u, v \in \{r, s, t\}$. Then equations (8) produce the following system of linear equations:

$$\begin{pmatrix} 1 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & d_t & -d_s & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & d_r & -d_t & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & d_s & -d_r \end{pmatrix} \begin{pmatrix} X_{rr} \\ X_{ss} \\ X_{tt} \\ X_{rs} \\ X_{rt} \\ X_{st} \\ X_{sr} \\ X_{tr} \\ X_{ts} \end{pmatrix} = \begin{pmatrix} d_r^2 - d_r \\ d_s^2 - d_s \\ d_t^2 - d_t \\ d_r d_s - T \\ d_t d_r - T \\ d_s d_t - T \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

This system has the unique solution from which we obtain the following expressions for the structure constants:

$$c_{u,u}^v = c_{u,u}^t = T \frac{d_u - d'_u}{2d_t d_v} + \frac{d_u}{2}, \quad c_{u,u}^u = T \frac{d'_u (d'_u - d_u)}{2m} + \frac{2d_u - 2 - d'_u}{2}, \quad (9)$$

where $\{u, v\} = \{r, s\}$ and $m = d_r d_s d_t$. By substituting these solutions to equality (3) with $u = r$ we obtain the following quadratic equation with respect to T :

$$(n-1)^2 T^2 - 2m(n+1)T + m^2 = 0.$$

Thus $T = m/(\sqrt{n} + \varepsilon_{r,s})^2$ where $\varepsilon_{r,s} = \pm 1$. (We note that due to condition (C3) we have $\sqrt{n} + \varepsilon_{r,s} > 0$.) Taking $u = r, s$ in the second of equalities (9) we see that in fact the number $\varepsilon_{r,s}$ does not depend on the choice of r, s .

Denote it by ε . Then by the definition of T this proves the first equality in (7); the other ones follow from (9) because due to Theorem 3.2 we have $c_{r,s}^w = c_{r,s}^t$ for all $w \in \mathcal{R}^\# \setminus \{r, s\}$. ■

Let us rewrite formulas (7) in slightly different form. For $r \in \mathcal{R}$ set $g_r = m + \varepsilon$ where $m = \sqrt{n}$. Then a straightforward computation shows that

$$d_r = g_r(m + \varepsilon), \quad c_{r,r}^r = (g_r + \varepsilon)(g_r + 2\varepsilon) - \varepsilon m - 2, \quad c_{r,r}^s = g_r(g_r + \varepsilon) \quad (10)$$

for all $s \in \mathcal{R} \setminus \{r\}$. Suppose from now that the C -algebra $\mathcal{A}$ is the adjacency algebra of an association scheme. Then its structure constants are nonnegative integers. This implies that the number n is an exact square and relations (10) show that each basis graph of the association scheme is a strongly regular graph of Latin square type (for $\varepsilon = -1$) or of negative Latin square type (for $\varepsilon = 1$). Thus as a special case of Theorem 3.3 we obtain the result of A. V. Ivanov who proved in [12] that the number of points of a nontrivial amorphic association scheme is an exact square and all basis graphs of the scheme are simultaneously strongly regular graphs of Latin square type or of negative Latin square type.

Theorem 3.3 enables us to compute the automorphism group of an arbitrary amorphic C -algebra $\mathcal{A} = (A, \mathcal{R})$. To do this denote by $\text{Aut}(D(\mathcal{A}))$ the group of all permutations $g \in \text{Sym}(\mathcal{R})$ such that $e^g = e$ and $d_r = d_{rg}$ for all $r \in \mathcal{R}^\#$. Clearly,

$$\text{Aut}(D(\mathcal{A})) = \prod_{C \in \pi_{\mathcal{A}}} \text{Sym}(C)$$

where $\pi_{\mathcal{A}}$ is the partition of $\mathcal{R}$ such that $C \in \pi_{\mathcal{A}}$ iff $C = \{e\}$ or $C = \{r \in \mathcal{R}^\# : d_r = d\}$ with $d \in \mathcal{D}(\mathcal{A})$.

Theorem 3.4 *Let $\mathcal{A} = (A, \mathcal{R})$ be a nontrivial amorphic C -algebra. Then $\text{Aut}(\mathcal{A}) = \text{Aut}(D(\mathcal{A}))$.*

Proof. The inclusion $\text{Aut}(\mathcal{A}) \leq \text{Aut}(D(\mathcal{A}))$ is obvious. To prove the converse inclusion it suffices to verify that

$$c_{rg,sg}^{tg} = c_{r,s}^t, \quad g \in \text{Sym}(D(\mathcal{A})), \quad r, s, t \in \mathcal{R}.$$

However, this follows from (8) because $d_{ug} = d_u$ for all $g \in \text{Aut}(D(\mathcal{A}))$. ■

3.3 In this subsection given a multiset D of at least 4 positive real numbers with $1 \in D$, and given an integer $\varepsilon \in \{-1, +1\}$ we construct the class $\mathcal{A}_\varepsilon(D)$ of all isomorphic nontrivial amorphic C -algebras. Let us start with the explicit construction.

Algebra $\mathcal{A}_\varepsilon(\mathcal{R}, D)$. Let $\mathcal{R}$ be a finite set with distinguished element e , and let $d : \mathcal{R} \mapsto D$ be a bijection from $\mathcal{R}$ to D such that $d_e = 1$. Denote by A the linear space over $\mathbb{C}$ spanned by $\mathcal{R}$ and by $*$ the identity map of A . Given $r, s, t \in \mathcal{R}$ set $c_{r,s}^t$ to be the real number defined by condition (C3) if $t = e$, by formula (7) with $n = \sum_{r \in \mathcal{R}} d_r$ if $e \notin \{r, s, t\}$ and $r \neq s \neq t \neq r$ or $r = s \neq t$ or $r = s = t$, and by equalities (4) otherwise. Then the linear space A with the multiplication defined by (1) becomes an (possibly non-associative) algebra over $\mathbb{C}$ in which the set $\mathcal{R}$ is a linear base.

Lemma 3.5 *The pair $\mathcal{A}_\varepsilon(\mathcal{R}, D) = (A, \mathcal{R})$ is a nontrivial amorphic C -algebra.*

Proof. Let us check that the algebra A is associative. To do this we need to verify that equalities (3) hold for all $r, s, t \in \mathcal{R}$. Clearly, it is true for $e \in \{r, s, t\}$ or $y = e$ or $r = s = t$. Thus it suffices to check the cases $r \neq s \neq t \neq r$ and $r = s \neq t$, and in each of them to verify separately all the possibilities for y : $y = r$ or $y = s$ or $y = t$ or $y \notin \{r, s, t\}$. All these computations are straightforward but too long to present here. For example, if $r \neq s \neq t \neq r$ and $y = s$, equality (3) is equivalent to the following relation:

$$c_{r,s}^r c_{r,t}^s + c_{r,s}^s c_{s,t}^s + c_{r,s}^t c_{t,t}^s = c_{r,r}^s c_{s,t}^r + c_{r,s}^s c_{s,t}^s + c_{r,t}^s c_{s,t}^t$$

which can be verified by the explicit substitution in it the values of $c_{r,s}^t$ given by (7) and (4).

Thus the algebra A is associative. It is easy to see that conditions (C1), (C2) and (C3) are satisfied. Finally, from the proof of Theorem 3.3 (see the first two lines of (8)) it follows that equalities (5) hold and hence condition (C4) is also satisfied. Thus $\mathcal{A}_\varepsilon(\mathcal{R}, D)$ is a C -algebra. The third line of (8) shows that the structure constants of A satisfies (6). So the C -algebra $\mathcal{A}_\varepsilon(\mathcal{R}, D)$ is amorphic by Theorem 3.2. ■

From Theorem 3.3 it follows that two C -algebras $\mathcal{A}_\varepsilon(\mathcal{R}, D)$ and $\mathcal{A}_{\varepsilon'}(\mathcal{R}', D')$ are isomorphic iff $\varepsilon = \varepsilon'$ and $D = D'$ (as multisets). Thus denoting by $\mathcal{A}_\varepsilon(D)$ the class of all C -algebras $\mathcal{A}_\varepsilon(\mathcal{R}, D)$ with fixed D and ε , we come to the following characterization of nontrivial amorphic C -algebras.

Theorem 3.6 *The class of all nontrivial amorphic C -algebras equals the union of classes $\mathcal{A}_\varepsilon(D)$ where D runs over multisets of at least four positive real numbers with $1 \in D$, and $\varepsilon \in \{-1, +1\}$. ■*

We complete the section by giving an example of amorphic C -algebra generalizing affine C -algebras.

Generalized affine C -algebras. Let $\mathcal{A} \in \mathcal{A}_\varepsilon(D)$ be a homogeneous C -algebra for some ε and D . Then the number d_r does not depend on the choice of the basis element r of $\mathcal{A}$ other than e . Denote it by d and set $n = n(\mathcal{A})$. We say that $\mathcal{A}$ is a *generalized affine C -algebra* if $d = \sqrt{n} + \varepsilon$. In this case from (7) it follows that

$$c_{r,s}^t = \begin{cases} 1, & \text{if } |\{r, s, t\}| = 3, \\ \varepsilon + 1, & \text{if } |\{r, s, t\}| = 2, \\ \varepsilon(3 - \sqrt{n}) + 1, & \text{if } |\{r, s, t\}| = 1 \end{cases} \quad (11)$$

for all basis elements r, s, t of $\mathcal{A}$ other than e . By comparing (2) and (11) with $(n, \varepsilon) = (q^2, -1)$ we see that any affine C -algebra corresponding to affine plane of order q , is a generalized affine C -algebra. The converse statement is not true because there are a lot of integers q for which there is no any affine plane of order q [7, p.63]

4 Homogeneous amorphic C -algebras

4.1 From the point of view of algebraic combinatorics C -algebras with non-negative integer structure constants are of special interest. The following result shows that in studying amorphic C -algebras of such a kind one can restrict himself to the homogeneous case. Since for a homogeneous C -algebra belonging to a class $\mathcal{A}_\varepsilon(D)$ we have $D = \{1, d, \dots, d\}$ where the positive real number d arises $\nu - 1$ times with ν being the dimension of the algebra, we write $\mathcal{A}_\varepsilon(\nu, d)$ for the class of all $\mathcal{A}_\varepsilon(\mathcal{R}, D)$ with $|\mathcal{R}| = \nu$.

Theorem 4.1 *Let $\mathcal{A} = \mathcal{A}_\varepsilon(\mathcal{R}, D)$ be an amorphic C -algebra for some ε , $\mathcal{R}$ and D . Suppose that a positive real number d has the following property:*

$$\{d_r/d : r \in \mathcal{R}^\#\} \subset \mathbb{N}. \quad (12)$$

Then $\mathcal{A}$ is a fusion of a homogeneous amorphic C -algebra belonging to the class $\mathcal{A}_\varepsilon(\nu, d)$ where $\nu = (n - 1)/d + 1$ with $n = n(\mathcal{A})$.

Proof. Set $\mathcal{A}' = \mathcal{A}_\varepsilon(\mathcal{R}', \mathcal{D}')$ where $\mathcal{R}'$ is a set of cardinality ν with a distinguished element e' and $\mathcal{D}' = \{1, d, \dots, d\}$ ($\nu - 1$ times). Then from Theorem 3.6 it follows that $\mathcal{A}'$ is a homogeneous amorphic C -algebra. So $\mathcal{A}' \in \mathcal{A}_\varepsilon(\nu, d)$. This implies that for any partition π' of $\mathcal{R}'$ such that $\{e'\} \in \pi'$ the π' -fusion of it is an amorphic C -algebra. Take $\pi' = \{C_r : r \in \mathcal{R}\}$ where C_r is a subset of $\mathcal{R}'$ of cardinality d_r/d . Then it is easy to see that $\mathcal{A}'_{\pi'} \in \mathcal{A}_\varepsilon(\mathcal{D})$. Taking into account that $\mathcal{A} \in \mathcal{A}_\varepsilon(D)$, we conclude that $\mathcal{A}$ is isomorphic to $\mathcal{A}'_{\pi'}$ as desired. ■

Corollary 4.2 *Any nontrivial amorphic C -algebra with integer structure constants is a fusion of a generalized affine C -algebra.*

Proof. Let $\mathcal{A}$ be a nontrivial amorphic C -algebra. Then by Theorem 3.6 $\mathcal{A} \in \mathcal{A}_\varepsilon(D)$ for some ε and D . Suppose that all the structure constants of $\mathcal{A}$ are integers. Then from the first line of (7) it follows that the numbers $d = \sqrt{n} + \varepsilon$ and d_r/d are integers for all basis elements $r \neq e$ of $\mathcal{A}$. By Theorem 4.1 this implies that $\mathcal{A}$ is a fusion of a homogeneous C -algebra $\mathcal{A}' \in \mathcal{A}_\varepsilon(\nu, d)$ where $\nu = (n - 1)/d + 1$ with $n = n(\mathcal{A})$. Since $n(\mathcal{A}') = n(\mathcal{A})$ it follows that $\mathcal{A}'$ is a generalized affine C -algebra. ■

The hypothesis of Theorem 4.1 is satisfied whenever the structure constants of the C -algebra $\mathcal{A}$ are rational numbers. Indeed, as the number d for which condition (12) holds one can choose any rational number $1/N$ where N is a multiple of the least common multiple of the denominators of the rational numbers d_r . Thus, any amorphic C -algebra with rational structure constants is a fusion of infinitely many nonisomorphic amorphic homogeneous C -algebras.

4.2 From Theorem 3.4 it follows that if $\mathcal{A}$ is a nontrivial homogeneous amorphic C -algebra, then $\text{Aut}(\mathcal{A}) \cong \text{Sym}(\mathcal{R}^\#)$. On the other hand, if $\mathcal{A} = (A, \mathcal{R})$ is an arbitrary $\mathbb{C}$ -algebra and $\pi = \text{Orb}(G, \mathcal{R})$ is a partition of $\mathcal{R}$ to the set of the orbits of a group $G \leq \text{Aut}(\mathcal{A})$, then the π -fusion of $\mathcal{A}$ is obviously a C -algebra. This implies the following statement.

Theorem 4.3 *A C -algebra $\mathcal{A} = (A, \mathcal{R})$ is homogeneous and amorphic iff $\text{Aut}(\mathcal{A}) \cong \text{Sym}(\mathcal{R}^\#)$. ■*

From the main result of the paper [11] it follows that the sufficiency in Theorem 4.3 can be strengthened for the case when the C -algebra $\mathcal{A}$ is the adjacency algebra of an association scheme. Namely, instead of the assumption

that $\text{Aut}(\mathcal{A}) \cong \text{Sym}(\mathcal{R}^\#)$ one can suppose that the group $\text{Aut}(\mathcal{A})$ has an elementary abelian 2-subgroup acting regularly on the set $\mathcal{R}^\#$. It seems to be true that this result does valid for arbitrary C -algebras (see [3]). It would be interesting to describe the properties of the group $\text{Aut}(\mathcal{A})$ providing that the C -algebra $\mathcal{A}$ is homogeneous and amorphic.

4.3 We recall that a *table algebra* is a C -algebra with nonnegative structure constants (see [1] and the references in it). In this subsection we deal with the homogeneous amorphic table algebras. In contrast to the general case this class of table algebras is selfdual (concerning the duality theory for C -algebras we refer to [4]). More exactly, the following statement holds.

Theorem 4.4 *Let $\mathcal{A}$ be a homogeneous amorphic table algebra and let $\hat{\mathcal{A}}$ be the dual C -algebra of $\mathcal{A}$. Then $\hat{\mathcal{A}}$ is also a homogeneous amorphic table algebra. Moreover, if $\mathcal{A} \in \mathcal{A}_\varepsilon(\nu, d)$ for some ε, ν, d , then $\hat{\mathcal{A}} \in \mathcal{A}_\varepsilon(\nu, d)$. In particular, the table algebras $\mathcal{A}$ and $\hat{\mathcal{A}}$ are isomorphic.*

Proof. Set $\mathcal{A} = (A, \mathcal{R})$ and $\hat{\mathcal{A}} = (\hat{A}, \hat{\mathcal{R}})$. Let $G \leq \text{Sym}(R)$ be a group acting regularly on $\mathcal{R}^\#$. Then $G \leq \text{Aut}(\mathcal{A})$ by Theorem 4.3. So by [11, Proposition 1.1]² there exists a group $\hat{G} \leq \text{Aut}(\hat{\mathcal{A}})$ such that $\hat{G} \cong G$ and $\hat{G}$ acts regularly on $\hat{\mathcal{R}}^\#$. When the group G runs over all the subgroups of $\text{Sym}(R)$ acting regularly on $\mathcal{R}^\#$, the group $\hat{G}$ runs over all the subgroups of $\text{Sym}(\hat{R})$ acting regularly on $\hat{\mathcal{R}}^\#$. Thus $\text{Aut}(\hat{\mathcal{A}}) \cong \text{Sym}(\hat{\mathcal{R}}^\#)$ and hence by Theorem 4.3 the C -algebra $\hat{\mathcal{A}}$ is homogeneous and amorphic. So by Theorem 3.6 we have

$$\mathcal{A} \in \mathcal{A}_\varepsilon(\nu, d), \quad \hat{\mathcal{A}} \in \mathcal{A}_{\hat{\varepsilon}}(\nu, \hat{d}) \quad (13)$$

where $\nu = |\mathcal{R}| = |\hat{\mathcal{R}}|$, $d = d_r$ for all $r \in \mathcal{R}^\#$, $\hat{d} = d_{\hat{r}}$ for all $\hat{r} \in \hat{\mathcal{R}}^\#$ and $\varepsilon, \hat{\varepsilon} \in \{-1, +1\}$. Moreover, from [4, (5.23)] it follows that

$$d(\nu - 1) + 1 = \sum_{r \in \mathcal{R}} d_r = \sum_{\hat{r} \in \hat{\mathcal{R}}} d_{\hat{r}} = \hat{d}(\nu - 1) + 1 \quad (14)$$

from which we conclude that $d = \hat{d}$. Denote by P and Q the first and the second eigenmatrices of the C -algebra $\mathcal{A}$ (the rows and the columns of the matrix P (resp. Q) are indexed by the elements of $\mathcal{R}$ and $\hat{\mathcal{R}}$ (resp. $\hat{\mathcal{R}}$ and

²This result is stated for adjacency algebras of association schemes, but in fact it is true for arbitrary C -algebra

$\mathcal{R}$) respectively). Then from [11, Theorem 1.2] it follows that there exists a bijection $\mathcal{R} \rightarrow \widehat{\mathcal{R}}, r \rightarrow \widehat{r}$ such that $P_{r,\widehat{s}} = P_{s,\widehat{r}}$ and $Q_{\widehat{r},s} = Q_{\widehat{s},r}$ for all $r, s \in \mathcal{R}$. Set $n = n(\mathcal{A}) = n(\widehat{\mathcal{A}})$. Then by Theorems 5.5 and 5.7 and equality (5.26) of [4] we have

$$\frac{Q_{\widehat{s},r}}{d_{\widehat{s}}} = \frac{\overline{P}_{r,\widehat{s}}}{d_r}, \quad c_{r,r}^r = \frac{d_r^2}{n} \sum_{\widehat{s} \in \widehat{\mathcal{R}}^\#} \frac{1}{d_{\widehat{s}}^2} Q_{\widehat{s},r} Q_{\widehat{s},r} \overline{Q}_{\widehat{s},r}, \quad c_{\widehat{s},\widehat{s}}^{\widehat{s}} = \frac{d_{\widehat{s}}^2}{n} \sum_{r \in \mathcal{R}^\#} \frac{1}{d_r^2} P_{r,\widehat{s}} P_{r,\widehat{s}} \overline{P}_{r,\widehat{s}}$$

for all $r, s \in \mathcal{R}^\#$, where the bar means the complex conjugation. Now the above equalities along with the fact that $d_r = d = \widehat{d} = d_{\widehat{s}}$ for all $r, s \in \mathcal{R}^\#$, imply by condition (C2) that $c_{\widehat{r},\widehat{r}}^{\widehat{r}} = c_{r,r}^r$ for all $r \in \mathcal{R}$. So from (7) it follows that $\varepsilon = \widehat{\varepsilon}$ and we are done by (13).■

Remark 4.5 *Theorem 4.4 shows that given a homogeneous amorphic table algebra $\mathcal{A}$, the pair of C -algebras $\mathcal{A}$ and $\widehat{\mathcal{A}}$ are in positive duality in the sense of [9].*

4.4 In this subsection we find conditions for parameters of a homogeneous amorphic C -algebra $\mathcal{A} = (A, \mathcal{R}) \in \mathcal{A}_\varepsilon(\nu, d)$ that guarantee it to be a table algebra. To do this we rewrite formulas (7) in terms of d, ε and $n = 1 + (\nu - 1)d$ as follows:

$$c_{r,s}^t = \begin{cases} \frac{d^2}{(\sqrt{n} + \varepsilon)^2}, & \text{if } |\{r, s, t\}| = 3, \\ \frac{d}{2} \left(1 + \frac{2d - n + 1}{(\sqrt{n} + \varepsilon)^2} \right), & \text{if } |\{r, s, t\}| = 2, \\ \frac{d - n + 1}{2} \left(1 + \frac{2d - n + 1}{(\sqrt{n} + \varepsilon)^2} \right) + d - 1, & \text{if } |\{r, s, t\}| = 1 \end{cases} \quad (15)$$

where $r, s, t \in \mathcal{R}^\#$. Then $\mathcal{A}$ is a table algebra iff the numbers $c_{r,s}^t$ are non-negative, which is equivalent to the two following inequalities:

$$d \geq \frac{n - 1 - Q}{2} = -(1 + \varepsilon\sqrt{n}), \quad (16)$$

$$2d^2 - 3d(n - 1 - Q) + (n - 1)^2 - (n + 1)Q \geq 0, \quad (17)$$

where $Q = (\sqrt{n} + \varepsilon)^2$. Denote by θ the right-hand side of (16) and by $d^\pm$ the roots of the quadratic polynomial on d in the left-hand side of (17). Then a straightforward computation shows that

$$d^\pm = \frac{1 + \varepsilon\sqrt{n}}{2} \left(-3\varepsilon \pm \sqrt{9 + 4\varepsilon\sqrt{n}} \right).$$

and the system of two inequalities (16) and (17) has two families of solutions:

$$\theta \leq d \leq d^-, \quad d \geq \max\{\theta, d^+\}.$$

These solutions give a necessary and sufficient condition for the C -algebra $\mathcal{A}$ to be an amorphic homogeneous table algebra.

References

- [1] Z. Arad, Harvey I. Blau, *On Table Algebras and Applications to Finite Group Theory*, J. Algebra, **138** (1991), 137–185.
- [2] Z. Arad, E. Fisman, M. Muzychuk, *Generalized table algebras*, Israel J. Math., **114** (1999) 29–60.
- [3] E. Bannai, *On a theorem of Ikuta, Ito and Munemasa*, European J. Combin., **13** (1992), 1–3.
- [4] E. Bannai, T. Ito, *Algebraic combinatorics. I*, Benjamin/Cummings, Menlo Park, CA, 1984.
- [5] L. D. Baumert, W. H. Mills, R. L. Ward, *Uniform cyclotomy*, J. Number Theory, **14** (1982), 67–82.
- [6] J. A. Davis, Q. Xiang, *Amorphic association schemes with negative latin square type graphs*, arXiv:math/0409401, (2004), 13 pp.
- [7] P. Dembowski, *Finite Geometries*, Springer, Berlin, 1968.
- [8] John D. Dixon, Brian Mortimer *Permutation groups*, Springer-Verlag New York, Inc. (1994).
- [9] S. A. Evdokimov, I. N. Ponomarenko, A. M. Vershik, *C-algebras and algebras in Plancherel duality*, Zapiski Nauchnykh Seminarov POMI, **240** (1997), 53–66. English translation: J. Math. Sci., New York, **96** (1999), 5, 3478–3485.
- [10] J. Y. Gol’fand, M. H. Klin *Amorphic cellular rings I*, in: *Investigations in Algebraic Theory of Combinatorial Objects*, VNIISI, Moscow, Institute for System Studies, 1985, 32–38(in Russian).

- [11] T. Ikuta, T. Ito, A. Munemasa, *On pseudo-automorphisms and fusions of an association scheme*, European J. Combin., **12** (1991), 317–325.
- [12] A. V. Ivanov, *Amorphic cellular rings II*, in: *Investigations in Algebraic Theory of Combinatorial Objects*, VNIISI, Moscow, Institute for System Studies, 1985, 39–49(in Russian).
- [13] T. Ito, A. Munemasa, M. Yamada, *Amorphous association schemes over the Galois rings of characteristic 4*, European J. Combin., **12** (1991), 513–526.